

Allergies: NKDA _____

NURSING INTERVENTIONS: (Line through to cancel orders)

- Prior to initiating therapy, assess patient for appropriate level of sedation
- Hold paralytic daily to facilitate physical/ neurological assessment. Resume infusion at prior dose if indicated by patient response
- SCDs
- Elevate Head of Bed
- Train-of-four (TOF) every shift, after dose changes, and PRN. Adjust infusion to obtain goal TOF 1:4 per protocol or physician order
- ROM every 4 hours **AND** PRN.

MEDICATIONS: (Check box to activate)

Begin infusion at lowest effective dose

<input type="checkbox"/> vecuronium: Loading dose 0.15 mg/ kg IV push, then infuse at 0.8 - 1.2 mcg/ kg/ minute Do not use in hepatic or renal dysfunction	Paralytic
<input type="checkbox"/> atracurium (TRACRIUM): Loading dose 0.5 mg/ kg IV push, then infuse at 2 - 15 mcg/ kg/ minute	
<input type="checkbox"/> rocuronium (ZEMURON): Loading dose: 0.6 mg/ kg IV push, then infuse at 0.3 - 1 mg/ kg/ hour	
<input checked="" type="checkbox"/> mineral oil/ petrolatum (LACRILUBE) ophthalmic ointment apply ¼ inch ribbon to each eye every 4 hours AND PRN	Dry Eyes/ Corneal Damage Prevention

Physician Signature

Date

Time