

Allergies: _____

Notify Physician if patient has:

Acute CVA, aortic dissection, brain metastasis, active internal bleeding, severe hypertension refractory to medical treatment, pregnancy, recent major surgery, CPR or trauma less than 14 days, severe liver disease, PT greater than 15 seconds or INR greater than 1.7, active peptic ulcer disease, traumatic intubation or puncture of a non-compressible vessel within 10 days

MEDICATIONS: (check box to activate)

Access Device	Dose per lumen	1 lumen	2 lumens	3 lumens
PICC, Subclavian, Jugular, Hohn, Midline, Hickman, Groshong	<input type="checkbox"/> Alteplase (t-PA) 2 mg/ 2 mL, sent in a 10 mL syringe X 1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Port-A-Cath	<input type="checkbox"/> Alteplase (t-PA) 3 mg/ 3 mL, sent in a 10 mL syringe X 1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

NURSING CARE:

- Notify IV Therapy Team to assess line and administer Alteplase
- After 30 - 60 minutes, assess patency of catheter by aspirating 2 - 3 mL with a 10 mL syringe. Discard aspirated fluid
- If the catheter is patent, aspirate alteplase and flush with 20 mL of 0.9% NS and resume IV or flush per *IV Therapy - Line Maintenance Protocol*
- If patency is not restored, notify Physician

Physician Signature

Date

Time



OVERLAKE Hospital Medical Center
Medical excellence every day™

DECLOTTING CENTRAL LINE ORDERS

P0094B (Rev 0909) White – Chart Yellow – Pharmacy Pink – Nursing