

ALLERGIES: _____

PRE – PROCEDURE ORDERS

Prep:

1. Notify Medical Imaging for Radiologist to obtain consent
2. Pre-menopausal female pts (12-57yr) to sign Pregnancy Consent Form
3. Start IV with 1000cc D5½ NS @ 100cc/hr. (For diabetic patients, substitute ½ NS)
4. NPO for 4 hours pre-procedure
5. Have Interventional Radiologist initial site of entry on patients going to OR.

Medications:

Pre-op on-call give:

- Diazepam (Valium) 5mg PO if age < 65
- Lorazepam (Ativan) 1 mg PO if age ≥ 65

Diagnostics:

- PT/INR
- CBC
- _____

POST-PROCEDURE ORDERS

Monitoring Patient:

1. Vital Signs/O₂ sat: q 30min x 2, q 1 hr X _____, q 4 hr/or per prior orders
2. NPO, patient going to OR
 Diet as tolerated when awake and alert
 Per prior orders
3. Activity as tolerated/or per prior orders
4. Run current IV at ____cc/hr for ____hr, then heparin lock until discharged
 IV per prior orders
5. I & O, measure nephrostomy/biliary catheter drainage and maintain outflow record.

Medications:

1. Acetaminophen(Tylenol) 325-650 mg PO q 4 hrs PRN fever/pain, **not to exceed 4g/24 hrs**

Nephrostomy/Biliary Tube Care:

1. Dressing changes as needed. (Call Interventional Radiologist if patient has increased drainage around catheter.)
2. Catheter to gravity drainage
3. Clamp nephrostomy/Biliary catheter at _____
(Place catheter back to gravity drainage if patient develops pain or fever. Notify IR at #5507)

Discharge Instructions:

- Pt may be discharged _____hrs post procedure if vital signs stable, fluid intake adequate, pain controlled, and without evidence of bleeding or hematoma.
- Discharge per Dr _____
- Give patient discharge instructions for Nephrostomy tube care.
- Schedule patient for follow-up nephrostogram/cholangiogram in General Radiology ext: 5507

MD Signature

Date

Time



NEPHROSTOMY/BILIARY TUBE PLACEMENT
Vascular & Interventional Radiology
P0115 (8/05)
White: Chart **Yellow:** Pharmacy