

Allergies: NKDA _____

Diagnosis: Morbid obesity

Procedure: Roux-EN-Y Gastric Bypass Adjustable Gastric Band Other: _____
 Open Laparoscopic Revision

Patient Weight _____Kg

Patient Height _____Inches

Patient Girth _____Inches

NURSING INTERVENTIONS: (Check box to activate. Line through pre-checked order to cancel)

- Fingerstick Blood Glucose on all diabetic patients & cover as directed below
- Fingerstick Blood Glucose & cover as directed below
- Baseline room air O₂ saturation with head of bed at 30 degrees

LABORATORY/ DIAGNOSTICS: (Check box/ fill in to activate)

- Chest X-Ray PA & Lateral, reason: _____
- ECG CBC with Diff Renal Panel PT/ INR Serum Pregnancy Test ABG

Blood: Type and Screen (determines Hemosafe eligibility). If not Hemosafe eligible, notify physician
(Hemosafe-ineligible means history of antibodies, so pre-ordering RBCs ensures blood is available on site. Hemosafe-eligible means blood is immediately available when transfusion is ordered.)

DIETARY: NPO

IV THERAPY: (Line through order to cancel)

- Lactated Ringers to infuse at 100 mL/ hour. If renal failure on dialysis, Normal Saline to infuse at 10 mL/ hour

MEDICATIONS: (Check box to activate. Line through pre-checked orders to cancel)

<input type="checkbox"/> ceFAZolin (ANCEF) 2 g IV X 1 preop. If allergic, clindamycin 900 mg IV X 1 <i>Antibiotics must be given within 60 minutes prior to incision</i>	Adjustable Gastric Band Prophylaxis Antibiotic
<input type="checkbox"/> ceFAZolin (ANCEF) 2 g IV X 1 preop AND metroNIDAZOLE (FLAGYL) 500 mg IV X 1 preop <input type="checkbox"/> cefOXitin (MEFOXIN) 2 g IV X 1 preop <input type="checkbox"/> clindamycin 900 mg IV X 1 preop AND aztreonam 2 g IV X 1 preop <i>Antibiotics must be given within 60 minutes prior to incision</i>	Stapled Bariatric Procedure Prophylaxis Antibiotic
<input type="checkbox"/> heparin 5000 units subcutaneously X 1, two hours before surgery AND SCDs or venous foot pumps if SCDs do not fit <input type="checkbox"/> dalteparin (FRAGMIN) 2500 units subcutaneously X 1, one to two hours before surgery AND SCDs or venous foot pumps if SCDs do not fit	VTE Prophylaxis
<input type="checkbox"/> SCDs or venous foot pumps if SCDs do not fit	VTE Prophylaxis with high risk for bleeding
Patients on a beta-blocker at home must have received a dose within 24 hours of surgery to meet SCIP requirements. Consider if cardiovascular disease, IDDM, creatinine greater than or equal to 2 <input type="checkbox"/> Beta Blocker: _____ (if not already taken) <input type="checkbox"/> metoprolol 12.5 mg PO X 1 preop if no beta blocker taken within previous 24 hours <input type="checkbox"/> Not indicated (e.g. asthma, recent bronchospasm, allergic, second/ third degree block)	Beta-blocker
<input checked="" type="checkbox"/> lispro (humaLOG) insulin subcutaneously X 1 dose according to the following blood glucose levels: 131-150 = 2 units; 151-180 = 4 units; 181-240 = 6 units; 241-300 = 8 units; 301-350 = 10 units; greater than 350 = 12 units & notify physician	Blood Glucose

Physician Signature

Date

Time

P H Y S I C I A N O R D E R



BARIATRIC SURGERY PREOP ORDERS
P0088D (Rev 0711) White – Chart/ Scan to Pharmacy