

ALLERGIES: _____

PRE-PROCEDURE

1. Clear liquids after midnight then NPO 3 hrs. pre-procedure.
2. IV: D₅ ½ NS @ 100cc/hr. If patient is diabetic, substitute ½ NS @ 100cc/hr.
3. Lab Work: INR CBC Coag panel Bleeding time for Cr > 4.0
 RP (if not done last 2 wks.)
4. Patient to void prior to procedure.
5. Medications to be given after consent signed:
 Diazepam (Valium) 5mg PO
 Lorazepam (Ativan) 1mg PO
 Diphenhydramine (Benadryl) 25-50mg PO for patients > 70 years old

6. Completed UW form goes with chart to Medical Imaging.

POST-PROCEDURE

1. VS Q 15 mins. x 4, then Q 30 mins. x 2; Q 1 hr. x 2; Q 2 hrs. x 2; then PRN until discharged. Check biopsy site for swelling, bleeding or discoloration when VS are taken.
2. Chest X-rays: Single PA view with inspiration immediately post biopsy. Have Interventional Radiologist read.
 Single PA view 3 hrs. post procedure.
3. Diet: As tolerated when awake and alert or cl liq x 4 hrs. then regular as tolerated.
4. Activity: Position patient so they are lying on puncture site for first hour.
 Bed rest ___ hrs. BRP after _____ hrs. May be in a semi-fowler position for eating.
 As tolerated.
5. IV Fluids: Run current IV at ___ cc/hr. for _____ hr., then saline lock until discharge.
 Saline lock IV until discharge.
6. Medications: Acetaminophen (Tylenol) 1-2 PO Q 4-6 hrs. PRN discomfort.
 Other: _____

**OBSERVE FOR PNEUMOTHORAX
OBSERVE FOR INTRA-ABDOMINAL HEMORRHAGE**

7. Notify Physician of any dyspnea, severe pain or significant changes in VS.
8. Discharge: Send patient home with Biopsy Discharge Instructions. Patient may be discharged if stable ___ hrs. post procedure.

MD Signature

Date

Time



BIOPSY ORDERS PRE / POST

P0036 (11/03)

White – Chart Yellow – Pharmacy Pink - Nursing